

Table 1. Distribution of Sensitivities of Clostridial Isolates to Four Antibiotics

Clostridium Species	Total Number	Penicillin (units/ml)						Aureomycin ($\mu\text{g}/\text{ml}$)						Terramycin ($\mu\text{g}/\text{ml}$)						Chloramphenicol ($\mu\text{g}/\text{ml}$)								
		.2	.4	.8	1.0	3.2	6.4	.0125	.025	.05	.1	.2	.8	.05	.1	.2	.4	.8	3.2	1	2	4	6	8	12	10	20	32
<i>Cl. bif fermentans</i>	17		2	2	10	3		1	2	12	2			1	11	3		2		2	4	5	3			3		
<i>Cl. butyricum</i>	1					1				1					1							1						
<i>Cl. lentoputrescens</i>	1	1								1				1						1								
<i>Cl. novyi</i>	8		2	2	1	1	2			2	3	3			1	4	2	1				4	1	1	1		1	
<i>Cl. paraputrificum</i>	2				2						1		1		1			1					2					
<i>Cl. perfringens</i>	7	1	2	2	2				1	1	3	2		1	1	1	3	1				1	2	2		2		
<i>Cl. sordelli</i>	2			1	1					1	1			1		1						1		3				1
<i>Cl. sporogenes</i>	11	1	3	2	2	1	2			3	7	1		1	6	2	1	1				1	3		2	1	1	
<i>Cl. tetanomorphum</i>	2		1	1						1	1			1		1						1	1					
<i>Cl. unclassified</i>	5		3		1		1			4		1		1	3	1					1	2	1			1		
Total	56	3	13	10	19	6	5	1	4	25	18	7	1	7	24	12	7	5	1	3	5	15	14	3	4	7	3	2