

**Table 3. Anaerobic Flora of Wounds and Blood, Series A, August-September 1952**

Clostridium Species	Source				Total Number
	Tissue*		Blood		
	Number	Strains (per cent)	Number	Strains (per cent)	
<i>Cl. sporogenes</i>	40	29.0	11	58	51
<i>Cl. perfringens</i>	29	21.0			29
<i>Cl. bifermentans</i>	15	10.9	4	21	19
<i>Cl. sordelli</i>	4	2.8			4
<i>Cl. multif fermentans</i>	11	8.0	1	5	12
<i>Cl. novyi</i>	10	7.2	3	16	13
<i>Cl. paraputrificum</i>	5	3.6			5
<i>Cl. tetanomorphum</i>	4	2.9			4
<i>Cl. butyricum</i>	3	2.2			3
<i>Cl. tertium</i>	3	2.2			3
<i>Cl. lentoputrescens</i>	3	2.2			3
<i>Cl. tetani</i>	3	2.2			3
<i>Cl. aerofaecidum</i>	2	1.4			2
<i>Cl. histolyticum</i>	1	0.7			1
<i>Cl. carnis</i>	1	0.7			1
<i>Cl. cochlearium</i>	1	0.7			1
<i>Cl. putrefaciens</i>	1	0.7			1
<i>Cl. sphenoides</i>	1	0.7			1
<i>Cl. capitovale</i>	1	0.7			1
<b>Total</b>	<b>138</b>		<b>19</b>		<b>157</b>

\*Includes embedded clothing and fragments.