

**Table 8. Anaerobic Flora of Wounds and Blood, Series B,  
December-February 1952-1953**

<i>Clostridium</i> Species	Tissues		Blood	
	Number of Strains	Per Cent of Total	Number of Strains	Per Cent of Total
<i>Cl. sporogenes</i>	25	19.6	2	50.0
<i>Cl. perfringens</i>	23	18.1	1	25.0
<i>Cl. novyi</i>	7	5.5	-----	-----
<i>Cl. bifermentans</i>	4	3.1	-----	-----
<i>Cl. sordelli</i>	4	3.1	-----	-----
<i>Cl. multifermentans</i>	3	2.3	-----	-----
<i>Cl. lentoputrescens</i>	6	4.7	1	25.0
<i>Cl. paraputrificum</i>	7	5.5		
<i>Cl. tertium</i>	4	3.1		
<i>Cl. tetani</i>	5	3.9		
<i>Cl. carnis</i>	5	3.9		
<i>Cl. capitovale</i>	4	3.1		
<i>Cl. tetanomorphum</i>	3	2.3		
<i>Cl. fallax</i>	2	1.5		
<i>Cl. cochlearium</i>	2	1.5		
<i>Cl. aerofotidum</i>	2	1.5		
<i>Cl. sphenoides</i>	2	1.5		
<i>Cl. difficile</i>	1	.8		
<i>Cl. butyricum</i>	1	.8		
<i>Cl. putrefaciens</i>	2	1.5		
unclassified or unidentified	15	11.8		
<b>Total</b>	<b>127</b>		<b>4</b>	