

Table 9. Incidence of Clostridial Species in Tissues of 43 Cases Showing Cultures Positive for Clostridia, Series B

<i>Clostridium</i> Species	Present in Num- ber of Cases	Strains Recovered from All Positive Cases	Per Cent Positive for Species	Tissues Cultured in Positive Cases	Per Cent of Positive Tissues
<i>Cl. perfringens</i>	13	23	30.2	78	29.4
<i>Cl. sporogenes</i>	13	25	30.2	69	36.2
<i>Cl. novyi</i>	5	7	11.5	26	26.9
<i>Cl. multifementans</i>	2	3	4.6	10	30.0
<i>Cl. bifementans</i>	3	4	6.9	17	23.5
<i>Cl. sordelli</i>	3	4	6.9	16	25.0
<i>Cl. lentoputrescens</i>	6	6	13.9	31	19.3
<i>Cl. paraputrificum</i>	7	7	16.2	31	22.5
<i>Cl. tetanomorphum</i>	2	3	4.6	7	42.8
<i>Cl. tertium</i>	4	4	6.9	13	30.7
<i>Cl. difficile</i>	1	1	2.3	2	50.0
<i>Cl. butyricum</i>	1	1	2.3	14	7.1
<i>Cl. putrefaciens</i>	2	2	4.6	8	25.0
<i>Cl. capitovale</i>	4	4	6.9	36	11.1
<i>Cl. fallax</i>	2	2	4.6	10	20.0
<i>Cl. tetani</i>	5	5	11.5	33	15.1
<i>Cl. cochlearium</i>	2	2	4.6	7	28.5
<i>Cl. aërofoetidum</i>	2	2	4.6	8	25.0
<i>Cl. sphenoides</i>	2	2	4.6	6	33.3
<i>Cl. carnis</i>	5	5	11.5	21	23.8
unclassified	12	15	27.9	75	20.0