

**Table 3. Distribution of Clostridial Species in Wounds, Skin, Clothing, and Soil**

Clostridium Species	Percentage of Clostridia				
	Summer 42 Wounds	Winter 112 Wounds	Skin Swabs 40 Samples	Clothing 35 Samples	Soil 21 Samples
<i>Cl. sporogenes</i>	45.2	8.9	10	14.2	30.9
<i>Cl. perfringens</i>	30.9	8.9	10	22.5	90.4
<i>Cl. novyi</i>	9.5	3.5	5	-----	4.7
<i>Cl. paraputrificum</i>	9.5	0.8	-----	-----	-----
<i>Cl. lentoputrescens</i>	4.7	3.5	-----	-----	9.5
<i>Cl. bifermentans</i>	23.8	1.7	-----	-----	23.8
<i>Cl. sordelli</i>	-----	2.6	-----	5.7	-----
<i>Cl. pasteurianum</i>	-----	-----	-----	-----	9.5
<i>Cl. carnis</i>	-----	3.5	-----	-----	4.7
<i>Cl. tetani</i>	-----	1.7	2.5	5.7	-----
<i>Cl. tertium</i>	9.5	1.7	2.5	-----	4.7
<i>Cl. capitovale</i>	-----	2.6	2.5	-----	-----
<i>Cl. fallax</i>	-----	0.8	-----	5.7	-----
<i>Cl. putrefasciens</i>	-----	0.8	2.5	-----	-----
<i>Cl. tetanomorphum</i>	7.1	-----	5	-----	9.5
<i>Cl. paratubulinum</i>	-----	-----	2.5	5.7	-----
<i>Cl. butyricum</i>	4.7	0.8	-----	11.4	23.8
<i>Cl. difficile</i>	-----	-----	-----	-----	4.7
<i>Cl. aerofoetidum</i>	2.3	-----	2.5	-----	19.4
<i>Cl. subterminale</i>	-----	-----	-----	-----	-----
<i>Cl. multif fermentans</i>	14.2	1.7	5	-----	14.2
<i>Cl. sphenoides</i>	2.3	0.8	-----	-----	4.7
<i>Cl. histolyticum</i>	2.3	1.7	-----	-----	-----
<i>Cl. cochlearium</i>	2.3	0.8	2.5	5.7	-----
<i>Cl. microaerophilus</i>	-----	-----	2.5	14.2	23.8
unclassified	-----	2.6	5	-----	-----