

**Table 8. Comparative Statistics for Case Fatality Rates of Abdominal Wounds by Organs Injured—Korea, 1952–1953, and World War II, 1942–1945\***

Viscus	KOREA				WORLD WAR II			
	Deaths, All Cases		Complicated† Per-centage Dying	Uncom- plicated Per-centage Dying	Deaths, All Cases*		Complicated† Per-centage Dying	Uncom- plicated Per-centage Dying
	Number	Per-centage			Number	Per-centage		
Colon	140	15.0	18.5	9.3	1,106	37	41	23
Jejunum and Ileum	134	13.4	16.4	3.0	1,168	30	36	14
Liver	102	15.6	20.6	9.0	829	27	39	10
Stomach	45	17.5	22.8	0	416	41	42	29
Kidney	65	25.4	29.1	0	427	35	38	16
Spleen	54	15.0	18.6	0	341	25	30	12
Rectum	22	18.1	16.0	18.7	155	30	42	14
Bladder	91	9.4	13.3	0	155	30	34	0
Duodenum	17	41.1	43.7	0	118	56	56	50
Pancreas	9	22.2	25.0	0	62	58	57	100
Gallbladder	33	0	0	0	53	30	30	0
Ureter	4	50.0	50.0	0	27	41	42	0

\*Second Auxiliary Surgical Group.<sup>2</sup>

†Additional organ damage.