

TABLE 16.—BLOOD CHANGES IN 69 SEVERELY WOUNDED MEN CLASSIFIED
ACCORDING TO DEGREE OF SHOCK
D. Severe Shock—16 Patients

Case No.	Blood Volume and Total Hemoglobin Determinations <i>Percent deviation from normal</i>						Blood Pressure at Time of Blood Volume Determination <i>mm. Hg</i>	Time after Wounding <i>hours</i>	Blood Sugar ¹ <i>mg. per 100 cc.</i>
	Blood Volume			Hemoglobin					
	From Observed Value	By Correction		From Observed Value	By Correction				
		A	B		A	B			
93	-11.7	-34.0	-30.3	+12.0	+2.6	+2.6	(0)	14½	—
100	-28.4	-69.0	-65.0	-44.1	-76.6	-76.6	80/50	4½	—
108	-25.5	-38.9	-30.0	-61.8	-64.1	-64.1	58/0	3½	372
110	-35.7	-37.6	-37.6	-45.7	-47.4	-47.4	(0)	¾	—
122	-34.4	-55.0	-48.2	-45.9	-58.4	-58.4	50/30	4	87
125	-21.3	-33.0	-29.1	-44.9	-52.7	-52.7	72/58	23	125
130	-31.3	-51.6	-40.5	-51.6	-62.0	-62.0	60/20	5	272
131	-29.8	-43.6	-39.0	-45.1	-54.3	-54.3	40/20	5¾	177
139	-19.3	-75.7	-23.3	-50.5	-70.7	-54.5	(0)	9	—
147	-6.5	-24.1	-14.3	-37.5	-42.7	-42.7	80/48	13¾	165
149	-2.3	-31.3	-9.8	-26.1	-32.7	-32.7	48/7	17¾	115
A-21	-46.7	-71.0	-54.8	-75.6	-83.8	-83.8	68/50	4¾	234
A-28	-9.2	-28.0	-9.2	-38.7	-38.7	-38.7	64/20	6¾	218
A-29	-28.2	-54.9	-37.1	-54.7	-63.6	-63.6	80/38	5¾	382
A-34	+12.8	-16.6	-8.2	(0)	-19.0	-19.0	80/68	26¾	125
A-38	-24.6	-58.3	-48.7	-42.0	-66.1	-66.1	(0)	?	133

¹ Blood sugar was determined in another patient in severe shock (Case 112) 11¾ hours after wounding and found to be 216 mg. per 100 cc. of blood.

² Case of crush injury.

(0): Not measurable.