

TABLE 61.—AVERAGE NONPROTEIN NITROGEN AND ELECTROLYTES OF PLASMA IN PATIENTS IN WHOM OLIGURIA DEVELOPED

Time of Determination	Plasma					Serum
	Nonprotein Nitrogen <i>mg. per 100 cc.</i>	CO ₂ Combining Power <i>mEq./L.</i>	Chlorides <i>mEq./L.</i>	Phosphate <i>mEq./L.</i>	Magnesium <i>mEq./L.</i>	Sodium <i>mEq./L.</i>
Preoperative	39.6±2.6 (21 cases)	22.8±1.4 (19 cases)	100.2±0.8 (20 cases)	3.7±0.3 (16 cases)	2.0±0.2 (10 cases)	138.6 (3 cases)
Postoperative Days						
First	59.9±5.8 (27 cases)	25.7±1.2 (18 cases)	98.1±0.9 (26 cases)	2.5±0.3 (21 cases)	1.4 (3 cases)	146.0 (3 cases)
Second to Fourth	99.6±11.4 (34 cases)	25.3±0.8 (29 cases)	96.4±1.7 (33 cases)	3.2±0.3 (26 cases)	2.4±0.4 (7 cases)	136.8±3.3 (6 cases)
Fifth to Seventh	131.8±29.3 (11 cases)	23.1±1.5 (8 cases)	92.3±2.4 (11 cases)	3.5±0.5 (9 cases)	3.5 (3 cases)	124.5 (2 cases)
Eighth to Tenth	139.1±50.2 (7 cases)	22.2±3.0 (6 cases)	87.4±4.3 (7 cases)	4.1±0.5 (4 cases)	5.2 (1 case)	136.5 (2 cases)
Eleventh to Fifteenth	140.4±42.8 (5 cases)	18.3±2.0 (4 cases)	99.8±5.2 (5 cases)	3.1±0.5 (5 cases)	2.3 (2 cases)	151.0 (2 cases)