

TABLE 62.—AVERAGE NONPROTEIN NITROGEN AND ELECTROLYTES OF PLASMA IN PATIENTS IN WHOM ANURIA DEVELOPED

Time of Determination	Plasma					Serum
	Nonprotein Nitrogen <i>mg. per 100 cc.</i>	CO ₂ Combin- ing Power <i>mEq./L.</i>	Chlorides <i>mEq./L.</i>	Phosphate <i>mEq./L.</i>	Magnesium <i>mEq./L.</i>	Sodium <i>mEq./L.</i>
Preoperative	34.7 (3 cases)	23.9 (3 cases)	98.7 (3 cases)	2.8 (3 cases)	1.8 (2 cases)	—
Postoperative Days						
First	75.6±4.7 (8 cases)	23.0±3.0 (6 cases)	95.7±2.4 (8 cases)	2.7±0.3 (7 cases)	1.7 (1 case)	143.3 (2 cases)
Second to Fourth	190.4±13.9 (20 cases)	23.2±1.3 (17 cases)	88.9±2.0 (20 cases)	4.3±0.5 (18 cases)	2.3±0.4 (5 cases)	141.2±2.7 (14 cases)
Fifth to Seventh	257.1±14.9 (12 cases)	20.0±1.6 (11 cases)	90.7±3.8 (11 cases)	4.5±0.1 (10 cases)	2.4±0.2 (4 cases)	142.8±6.9 (5 cases)
Eighth to Tenth	258.0±21.8 (4 cases)	19.1±2.3 (4 cases)	76.7±4.2 (4 cases)	5.9±0.9 (4 cases)	1.8 (1 case)	138.8 (3 cases)
Eleventh to Fifteenth	155.0 (2 cases)	22.9 (2 cases)	92.0 (2 cases)	3.9 (2 cases)	1.8 (1 case)	148.1 (2 cases)