

TABLE 75.—AVERAGE SPECIFIC GRAVITY OF THE URINE IN PATIENTS WITH HIGH AZOTEMIA\*

Classification of Cases	Average Specific Gravity					
	Preoperative Period	Postoperative Period				
		1 Day	2-4 Days	5-7 Days	8-10 Days	11-15 Days
ALL CASES . . . . .	1.026±.002 (11 cases)	1.020±.001 (21 cases)	1.015±.001 (46 cases)	1.012±.001 (28 cases)	1.014±.001 (13 cases)	1.011±.001 (12 cases)
<i>Patients' Outcome</i>						
Lived . . . . .	1.026±.001 (4 cases)	1.020±.001 (5 cases)	1.015±.001 (15 cases)	1.012±.001 (13 cases)	1.012±.002 (7 cases)	1.011±.001 (10 cases)
Died—All Causes . . . . .	1.026±.003 (7 cases)	1.020±.002 (16 cases)	1.015±.001 (31 cases)	1.013±.001 (15 cases)	1.016±.002 (6 cases)	1.011 (2 cases)
Died of Uremia . . . . .	1.033 (2 cases)	1.021±.002 (9 cases)	1.014±.001 (26 cases)	1.011±.001 (11 cases)	1.013 (4 cases)	1.011 (2 cases)
Died of other Causes . . . . .	1.023±.003 (5 cases)	1.018±.002 (7 cases)	1.019±.002 (5 cases)	1.018±.003 (4 cases)	1.022 (2 cases)	—
<i>Urinary Output</i>						
Normal . . . . .	1.028 (1 case)	1.019 (2 cases)	1.016±.001 (7 cases)	1.016±.002 (8 cases)	1.017±.004 (4 cases)	1.007 (1 case)
Anuria . . . . .	1.032 (1 case)	1.019 (7 cases)	1.014 (18 cases)	1.010 (10 cases)	1.012 (3 cases)	1.013 (3 cases)
Oliguria . . . . .	1.026 (7 cases)	1.021 (12 cases)	1.016 (21 cases)	1.012 (10 cases)	1.013 (6 cases)	1.011 (6 cases)
Unknown . . . . .	1.020 (2 cases)	—	—	—	—	1.014 (2 cases)
Anuria and Oliguria . . . . .	1.027±.002 (8 cases)	1.020±.001 (19 cases)	1.015±.001 (39 cases)	1.011±.001 (20 cases)	1.013±.001 (9 cases)	1.011±.001 (9 cases)
Anuria, Oliguria, and Unknown . . . . .	1.026±.002 (10 cases)	—	—	—	—	1.012±.001 (11 cases)

\* If more than 1 determination had been made on 1 patient during the specified period, the lowest value was used.