

TABLE 76.—AVERAGE pH OF THE URINE IN PATIENTS WITH HIGH AZOTEMIA\*

Classification of Cases	Average pH of Urine					
	Preoperative Period	Postoperative Period				
		1 Day	2-4 Days	5-7 Days	8-10 Days	11-15 Days
ALL CASES . . . . .	5.8±0.2 (12 cases)	6.0±0.1 (21 cases)	6.2±0.1 (43 cases)	6.1±0.1 (29 cases)	6.0±0.1 (14 cases)	6.3±0.3 (12 cases)
<i>Patients' Outcome</i>						
Lived . . . . .	5.6±0.2 (4 cases)	6.2±0.3 (5 cases)	6.6±0.2 (15 cases)	5.2±0.3 (13 cases)	5.8±0.2 (7 cases)	6.4±0.3 (10 cases)
Died—All Causes . . . . .	5.9±0.2 (8 cases)	6.0±0.1 (16 cases)	5.9±0.1 (28 cases)	6.2±0.2 (16 cases)	6.2±0.2 (7 cases)	6.0 (2 cases)
Died of Uremia . . . . .	5.5 (2 cases)	5.9±0.2 (9 cases)	5.9±0.1 (23 cases)	6.4±0.3 (12 cases)	6.4±0.3 (5 cases)	6.0 (2 cases)
Died of other Causes . . . . .	6.0±0.3 (6 cases)	6.2±0.2 (7 cases)	6.1±0.3 (5 cases)	5.5±0.2 (4 cases)	5.7 (2 cases)	—
<i>Urinary Output</i>						
Normal . . . . .	5.7 (2 cases)	6.6 (2 cases)	6.2±0.2 (7 cases)	5.8±0.2 (8 cases)	6.0±0.2 (4 cases)	5.9 (1 case)
Anuria . . . . .	5.5 (1 case)	5.8 (7 cases)	5.9 (16 cases)	6.2 (11 cases)	6.4 (4 cases)	6.9 (2 cases)
Oliguria . . . . .	5.8 (7 cases)	6.1 (12 cases)	6.4 (20 cases)	6.1 (10 cases)	5.8 (6 cases)	5.9 (6 cases)
Unknown . . . . .	6.1 (2 cases)	—	—	—	—	6.9 (2 cases)
Anuria and Oliguria . . . . .	5.7±0.2 (8 cases)	6.0±0.1 (19 cases)	6.2±0.1 (36 cases)	6.2±0.2 (21 cases)	6.0±0.2 (10 cases)	6.2±0.3 (8 cases)
Anuria, Oliguria, and Unknown . . . . .	5.8±0.2 (10 cases)	—	—	—	—	6.3±0.3 (10 cases)

\* If more than 1 determination had been made on 1 patient during the specified period, the lowest value was used.