

TABLE 86.—MEAN PLASMA “HEMOGLOBIN” LEVELS IN RELATION TO SHOCK AND QUANTITY OF TRANSFUSED BLOOD—76 CASES

Quantity of Blood Received (units)	Plasma Hemoglobin Levels ¹					
	Blood Specimen Drawn Less than 24 Hours after Injury ²			Blood Specimen Drawn 24-96 Hours after Injury ²		
	Shock		Total Cases	Shock		Total Cases
	Minimal	Moderate-Severe		Minimal	Moderate-Severe	
None	12.4 (7 cases)	9.8 (7 cases)	11.1 (14 cases) 9.2 (5 cases)	9.2 (5 cases)	
1 to 3	14.0 (2 cases)	8.7 (20 cases)	9.2 (22 cases)	16.0 (10 cases)	12.4 (3 cases) 15.1 (13 cases)	
4 to 14	6.8 (2 cases)	7.9 (3 cases)	7.5 (5 cases) 20.5 (17 cases)	20.5 (17 cases)	
Average	11.7 (11 cases)	8.8 (30 cases)	9.6 (41 cases)	16.0 (10 cases)	17.3 (25 cases) 16.9 (35 cases)	

¹ Mg. of benzidine-reacting material per 100 cc. of plasma.

² Determinations based on first blood specimen obtained for this analysis.