

TABLE 2.—Results of treatment in 30 cases of proved kala-azar

Case No.	Treatment	Fuadin (cc.)	Tartar emetic (gm.)	Neostam (gm.)	Neostibosan (gm.)	Stibanose (cc.)	Stilbamidine (gm.)
1	I.V.				{ ¹ 3.5		
2	do.				{ ² 5.0		
					{ ² 2.9		
3	do.			{ ¹ 3.6	{ ² 5.0		
				{ ¹ 4.25			
4	do.					² 200	
5	do.			² 4.05			
6	I.M.			¹ 3.5	² 3.9		
7	do.	³ 30		¹ 2.9	² 3.9		
8	do.			² 3.5			
9	do.				² 3.6		
10	I.V.				² 5.1		
11	I.M.				² 3.6		
12	do.			² 3.9			
13	do.				² 4.7		
14	I.V.			⁴ 6.4	⁴ 2.9		
15	I.M.			² 3.9			
16	do.				² 3.9		
17	I.V.			¹ 4.7	{ ¹ 2.4		
					{ ² 5.0		
18	I.M.			¹ 4.0	² 5.0		
					{ ¹ 2.7		
19	I.V.	³ 43		³ 8	{ ¹ 5.0	¹ 240	² 4.0
					{ ¹ 10.0		
20	do.			¹ 4.6	² 5.0		
21	I.M.			¹ 4.5		² 200	
22	do.				² 5.0		
23	do.					¹ 200	² 4.0
24	I.V.			² 3.8			
25	I.M.		⁴ 1.9		⁴ 2.6		
26	I.V.			² 3.8			
27	do.			⁴ 2.6	⁴ 2.6		
28	I.M.			¹ 2.5	² 5.0		
29	I.V.				² 3.4		
30	do.				{ ¹ 3.4		
					{ ² 5.0		

¹ Clinical improvement followed by relapse.² Amount which produced cure.³ No effect.⁴ Continuous course of two drugs which produced cure.

NOTE.—I.V. (intravenous injection); I.M. (intramuscular injection).