

TABLE 15.—*Admissions and deaths from fever of undetermined origin<sup>1</sup> in the total U.S. Army and in the Latin American area, by year, 1942-45*

[Admission rate expressed as number per annum per 1,000 average strength]

[Death rate expressed as number per annum per 100,000 average strength]

| Year       | Total Army |       | Latin America |        |
|------------|------------|-------|---------------|--------|
|            | Number     | Rate  | Number        | Rate   |
| Admissions |            |       |               |        |
| 1942.....  | 7, 711     | 2. 38 | 4, 385        | 43. 02 |
| 1943.....  | 40, 163    | 5. 85 | 2, 340        | 19. 37 |
| 1944.....  | 52, 586    | 6. 75 | 951           | 11. 08 |
| 1945.....  | 42, 270    | 5. 69 | 1, 000        | 13. 95 |
| Total..... | 142, 730   | 5. 63 | 8, 676        | 22. 82 |
| Deaths     |            |       |               |        |
| 1942.....  | 3          | 0. 09 | -----         | 0      |
| 1943.....  | 5          | . 07  | -----         | 0      |
| 1944.....  | 13         | . 17  | -----         | 0      |
| 1945.....  | 5          | . 07  | -----         | 0      |
| Total..... | 26         | 0. 10 | -----         | 0      |

<sup>1</sup> Probably malaria in many cases.